

AWARD NUMBER: W81XWH-13-1-0075

TITLE: LincRNAs and AR Reactivation after Androgen Deprivation in Prostate Cancer Cells

PRINCIPAL INVESTIGATOR: Xian-Ming Chen, MD

CONTRACTING ORGANIZATION: Creighton University
Omaha, NE 68178-0002

REPORT DATE: December 2014

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE December 2014		2. REPORT TYPE Final		3. DATES COVERED 30 Sep 2013 - 29 Sep 2014	
4. TITLE AND SUBTITLE LincRNAs and AR Reactivation after Androgen Deprivation in Prostate Cancer Cells				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-13-1-0075	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Xian-Ming Chen, M.D. E-Mail:xianmingchen@creighton.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Creighton University 2500 California Plaza Omaha, NE 68178-0002				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Prostate cancer (PCa) represents the most frequently diagnosed malignancy of men in the US. Reactivation of androgen receptor (AR) signaling following androgen deprivation is a major driver of the development of castration-resistant prostate cancer (CRPC). Understanding the precise mechanisms underlying aberrant AR-regulated gene activation will advance our knowledge of PCa tumorigenesis, with implementation of new therapeutic strategies. LincRNAs are recently identified novel genetic materials. The role of linRNAs in cancer is being increasingly accepted, both as possible specific biomarkers and as potential therapeutic targets. However, how aberrant expression of these lincRNAs contributes to PCa progression is still not fully understood. This one-year hypothesis-exploratory project was designed to test a novel hypothesis that lincRNAs promote AR reactivation in prostate cancer cells following androgen deprivation through modulation of AR/EZH2-mediated repressive chromatin remodeling. The experiments as proposed were successfully completed during the funding period and objectives achieved. Resultant data provide a basis for the identification of novel targets for therapeutic intervention for PCa. The ultimate goal is to define the molecular mechanisms of PCa initiation and development and to develop effective treatments.					
15. SUBJECT TERMS Prostate Cancer; Tumor progression; lincRNAs; PCATs; Androgen receptor; Castration resistance; EZH2; Histone Modifications					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Unclassified	18. NUMBER OF PAGES 9	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
1. Introduction.....	1
2. Keywords.....	1
3. Overall Project Summary.....	1
4. Key Research Accomplishments.....	5
5. Conclusion.....	5
6. Publications, Abstracts, and Presentations.....	5
7. Inventions, Patents and Licenses.....	5
8. Reportable Outcomes.....	5
9. Other Achievements.....	5
10. References.....	5
11. Appendices.....	6

1. Introduction

Prostate cancer (PCa) represents the most frequently diagnosed malignancy of men in the US, with a lifetime incidence risk of 1 in 6, and an expected 241,000 new cases and 33,700 death in 2011. The androgen receptor (AR) is a member of a nuclear receptor superfamily that regulates gene transcription. Many of these AR-regulated genes are key regulators to prostate development and progression. Understanding the precise mechanisms underlying aberrant AR-regulated gene activation will advance our knowledge of PCa tumorigenesis, with implementation of new therapeutic strategies. LincRNAs are recently identified novel genetic materials. The human genome encodes thousands of lincRNAs and each of these molecules can control hundreds of genes^[1]. Therefore, these molecules are extremely powerful molecules that regulate almost all aspects of the host's well-being. Recent research indicates that many lincRNAs have been linked to the initiation and progression of human cancer. The role of lincRNAs in cancer is being increasingly accepted, both as possible specific biomarkers and as potential therapeutic targets. A subset of lincRNAs, so-called prostate cancer-associated ncRNA transcripts (PCATs), may play an active role in prostate cancer progression^[2]. However, how aberrant expression of these lincRNAs/PCATs contributes to PCa progression is still not fully understood. Reactivation of androgen receptor (AR) signaling following androgen deprivation is a major driver of the development of castration-resistant prostate cancer (CRPC). This one-year hypothesis-exploratory project was designed to test a novel hypothesis that lincRNAs/PCATs promote AR reactivation in prostate cancer cells following androgen deprivation through modulation of AR/EZH2-mediated repressive chromatin remodeling. Resultant data provide a basis for the identification of novel targets for therapeutic intervention for PCa. The ultimate goal is to define the molecular mechanisms of PCa initiation and development and to develop effective treatments for this critically important disease.

2. Keywords

Prostate Cancer; Tumor progression; lincRNAs; PCATs; Androgen receptor; Castration resistance; EZH2; Histone Modifications; NF- κ B; IKK α ; Chromatin immunoprecipitation; RNA immunoprecipitation.

3. Overall Project Summary

Hypothesis and Aims: AR has been known as a transcriptional activator inducing prostate-specific gene expression. Data from integrative genome-wide analysis indicate that AR also acts as a transcriptional repressor. This repression, dictated by EZH2-associated repressive chromatin remodeling such as histone modifications, plays an important role in prostate cancer progression^[3]. EZH2 is an RNA-binding protein and can interact with lincRNAs to induce gene repression through H3K27me3. Indeed, EZH2-lincRNA interactions have been demonstrated to direct complex patterns of chromatin remodeling complex, resulting in gene suppression in prostate cancer cells^[4]. Nevertheless, the role of lincRNA/EZH2-mediated chromatin remodeling in AR reactivation following androgen deprivation has not been investigated. Although prostate cancer is a heterogeneous disease and its progression involves many pathogenic factors, inflammatory responses elicited by androgen deprivation contribute to the emergence of CRPC^[5]. The nuclear factor kappa B (NF- κ B) signaling

pathway is the master regulator for inflammatory reactions, and CRPC cells usually exhibit activated NF- κ B. Many stimuli activate NF- κ B, mostly through activation of IKK α and IKK β in the cytoplasm, resulting in nuclear translocation of NF- κ B subunits and, subsequently, transactivation and transrepression of targeted genes. CRPC cells are usually with the accumulation of activated IKK α in nuclei. The kinase activity of nuclear IKK α mediates phosphorylation of regulatory proteins and is required for IKK α -mediated gene suppression^[5]. Intriguingly, phosphorylation of EZH2 can facilitate lincRNA binding to EZH2 and, consequently, chromatin remodeling^[6]. Therefore, the **overriding hypothesis** of this exploratory project is that nuclear accumulation of IKK α enhances lincRNA/PCATs-guided chromatin recruitment of EZH2 complex and EZH2/AR-mediated H3K27me3 modification, promoting AR reactivation in prostate cancer cells following androgen deprivation (Figure 1). Two specific aims were proposed to address this central hypothesis: **Aim 1** was designed to define that nuclear translocation of IKK α enhances lincRNA-guided chromatin recruitment of EZH2 complex in prostate cells. **Aim 2** were proposed to clarify that IKK α /lincRNA-induced chromatin recruitment of EZH2 promotes EZH2/AR-mediated H3K27me3 modification and contributes to the reactivation of AR signaling following androgen deprivation in prostate cancer cells.









Fig. 1. Our hypothesis on lincRNA/IKK α -mediated gene suppression associated with PCa progression. Aberrant activation of NF- κ B signaling will increase expression of NF- κ B-responsive lincRNAs and cause nuclear translocation of IKK α . Nuclear translocation of IKK α will promote the chromatin recruitment of PRC2/EZH2/lincRNA complex, contributing to the suppression of metastatic suppressor genes key to PCa progression.

Studies, Results and Significance: During the one-year funding period, we have actively addressed all the key components as proposed, using an array of biochemical and molecular biological approaches, such as chromatin immunoprecipitation (ChIP) analysis to identify chromatin recruitment of EZH2, RIP assay for EZH2-lincRNA interactions, H3K27me3 analysis, and RNA interference for functional manipulation. We have completed all the experiments as proposed in the SOW for both Specific Aims and have not encountered insurmountable technical problems. There were no important modifications to our original plans. Resultant data provide mechanistic insights into the molecular mechanisms that link chronic inflammation to AR reactivation in prostate cancer cells following androgen deprivation.

Specifically, we have screened the expression of lincRNAs in metastatic PCa tissues and cell lines, as well as in PCa cells at the androgen deprivation condition. A panel of NF- κ B-responsive lincRNAs was identified to be consistently upregulated in metastatic PCA tissues and cell lines, or PCa cells after androgen deprivation (Figure 2A) and expression levels of selected lincRNAs of the panel by PCR were shown in Figure 2B. These NF- κ B-responsive lincRNAs were selected according to the NF- κ B binding sites in their gene promoter regions

(Figure 2A). Importantly, induction of these lincRNAs was also detected in LNCaP cells after stimulation with lymphotoxin B (Figure 3). Lymphotoxin B is an activator to stimulate NF- κ B signaling and induce nuclear translocation of IKK α in PCa cells and thus, has been used as a positive control in our analysis. Consistent with the concept that aberrant activation of NF- κ B signaling in PCa progression, we detected a significant increase of nuclear translocation of IKK α in LNCaP cells at the androgen deprivation condition (Figure 4A). Accordingly, we also detected an increased assembly of a panel of NF- κ B-responsive lincRNAs to the PRC2/EZH2 complex in PCa cells after androgen deprivation by RIP analysis. Data showing the assembly of PCAT-1 and Hotair to the PRC2/EZH2 complex in LNCaP cells after androgen deprivation were shown in Figure 4B and Figure 4C, respectively. To identify the target genes that may be associated with PRC2/EZH2 promoter recruitment in PCa cells following androgen deprivation, we measured the expression level of those metastatic suppressive genes. We focused on those genes that have the ORC2-repression signature in their gene loci. Our analysis revealed several of those genes that are downregulated in PCa cells following androgen deprivation (Figure 5A). Downregulation of representative genes in LNCaP cells following androgen deprivation as measured by PCR was shown in Figure 5B. To further test the potential recruitment of PRC2/EZH2 complex and its association with suppressive histone modification, we performed ChIP analysis in LNCaP cells following androgen deprivation and our data demonstrate the recruitment of PRC2/EZH2/lincRNA complex to the gene loci that are downregulated in cells following androgen deprivation. As a representative example shown in Figure 6, using an antibody to EZH2 for the ChIP analysis, we detected an increase in PRC2/EZH2 complex to the KCNA1 gene promoter region. Accordingly, an enrichment of suppressive H3K27me3 was also detected in the nearby region of the KCNA1 gene promoter by ChIP analysis using an antibody against H3K27me3. In addition, we have also measured the involvement of EZH2-lincRNA interactions in EZH2/AR-mediated H3K27me3 modification in LNCaP cells in response to AR activation. All these data are currently in preparation for publication.





4. Key Research Accomplishments

- Demonstrating that IKK α induces phosphorylation of EZH2 and enhances lincRNA-guided chromatin recruitment of EZH2 complex in response to AR activation in prostate cells.
- Clarification that IKK α /lincRNA-induced chromatin recruitment of EZH2 promotes EZH2/AR-mediated H3K27me3 modification and contributes to the reactivation of AR signaling following androgen deprivation in prostate cancer cells.

5. Conclusion

In conclusion, this one-year hypothesis exploratory proposal was designed to address the DOD PCRP 2012 focus areas on Mechanisms of Resistance (Understanding primary and acquired resistance to therapy) and Tumor and Microenvironment Biology (Understanding prognosis and progression of prostate cancer). We have actively addressed all the key components and have completed all the experiments as proposed in the SOW during the funding period. Resultant data provide not only mechanistic insights into the molecular mechanisms that link chronic inflammation to AR reactivation in prostate cancer cells following androgen deprivation but also a basis for the identification of novel targets for therapeutic intervention for CRPC, addressing the overarching challenge of the DOD PCRP to develop effective treatments for advanced prostate cancer.

6. Publications, Abstracts, and Presentations

Resultant data are in preparation for publication.

7. Inventions, Patents and Licenses

(None).

8. Reportable Outcomes

(None).

9. Other Achievements

(None).

10. References

- 1) Guttman M, Amit I, Garber M, et al. Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature*. 2009;458:223-7.
- 2) Prensner JR, Iyer MK, Balbin OA, et al. Transcriptome sequencing across a prostate cancer cohort identifies PCAT-1, an unannotated lincRNA implicated in disease progression. *Nat Biotechnol*. 2011;29:742-9.

- 3) Zhao JC, Yu J, Runkle C, et al. Cooperation between Polycomb and androgen receptor during oncogenic transformation. *Genome Res.* 2012;22:322-31.
- 4) Crea F, Hurt EM, Mathews LA, et al. Pharmacologic disruption of Polycomb Repressive Complex 2 inhibits tumorigenicity and tumor progression in prostate cancer. *Mol Cancer.* 2011;10:40.
- 5) Kaneko S, Li G, Son J, et al. Phosphorylation of the PRC2 component Ezh2 is cell cycle-regulated and up-regulates its binding to ncRNA. *Genes Dev.* 2010;24:2615-20.
- 6) Ammirante M, Luo JL, Grivennikov S, Nedospasov S, Karin M. B-cell-derived lymphotoxin promotes castration-resistant prostate cancer. *Nature.* 2010;464:302-5.

11. Appendices

(None).